
TOP 5 THINGS EVERYONE NEEDS TO KNOW ABOUT





Zika primarily spreads through infected mosquitoes. You can also get Zika through sex.

Many areas in the United States have the type of mosquitoes that can spread Zika virus. One type of mosquito that could spread Zika has been captured in Fairfield County in past years. Over years of testing, no mosquitoes that spread Zika have been found in Glastonbury. These mosquitoes are aggressive daytime biters and can also bite at night. Also, a man infected with Zika can pass it to others through sex.

- The best way to prevent Zika and other mosquito-transmitted diseases is to prevent mosquito bites.
 - Use insect repellent. It works!
 - Wear long-sleeved shirts and long pants.
 - Stay in places with air conditioning or window and door screens.
 - Remove standing water around your home.



3




Zika is linked to birth defects.

Zika infection during pregnancy can cause a serious birth defect called microcephaly that is a sign of incomplete brain development. If you have a male partner who lives in or has traveled to an area with Zika, do not have sex, or use condoms the right way, every time, during your pregnancy.

Pregnant women should not travel to areas with Zika.

If you must travel to one of these areas, talk to your healthcare provider first and strictly followsteps to prevent mosquito bites during your trip.





Returning travelers infected with Zika can spread the virus through mosquito bites.

Zika virus can stay in your blood for a week. When a mosquito bites you, you can pass the virus to the mosquito. The infected mosquito bites other people, who get infected.

